INSTITUT NATIONAL DE LA STATISTIQUE ET DES ETUDES ECONOMIQUES Série des Documents de Travail du CREST (Centre de Recherche en Economie et Statistique)

# n° 2003-33

# Sensitivity Analysis of Var and Expected Shortfall for Portfolios under Netting Agreements

J.-D. FERMANIAN<sup>1</sup> O. SCAILLET<sup>2</sup>

Les documents de travail ne reflètent pas la position de l'INSEE et n'engagent que leurs auteurs.

Working papers do not reflect the position of INSEE but only the views of the authors.

<sup>&</sup>lt;sup>1</sup> CDC Ixis Capital Markets, 47 quai d'Austerlitz, 75648 Paris Cedex 13 and CREST, 15 boulevard Gabriel Péri, 92245 Malakoff Cedex, France. Mail : <u>jdfermanian@cdcixis-cm.com</u>

<sup>&</sup>lt;sup>2</sup> HEC Genève and FAME, Université de Genève, 102 Bd Carl Vogt, CH-1211 Genève 4, Suisse. Mail : scaillet@hec.unige.ch

We thank G.P. Szegö and the referee for constructive criticism. We are grateful to Aubry Miens for having provided us the database. The second author receives support by the Swiss National Science Foundation through the National Center of Competence : Financial Valuation and Risk Management (NCCR FINRISK). Part of this research was done when he was visiting THEMA and IRES. Downloadable at http://www.hec.unige.ch/professeurs/SCAILLET Olivier/pages web/Home Page.htm

# SENSITIVITY ANALYSIS OF VAR AND EXPECTED SHORTFALL FOR PORTFOLIOS UNDER NETTING AGREEMENTS

Jean-David FERMANIAN<sup>a</sup> and Olivier SCAILLET<sup>b 1</sup>

<sup>a</sup> CDC Ixis Capital Markets, 47 quai d'Austerlitz, 75648 Paris cedex 13, and CREST, 15 bd Gabriel Péri,
 92245 Malakoff cedex, France. jdfermanian@cdcixis-cm.com

<sup>b</sup> HEC Genève and FAME, Université de Genève, 102 Bd Carl Vogt, CH - 1211 Genève 4, Suisse. scaillet@hec.unige.ch

<sup>1</sup>We thank G.P. Szegö and the referee for constructive criticism. We are grateful to Aubry Miens for having provided us the database. The second author receives support by the Swiss National Science Foundation through the National Center of Competence: Financial Valuation and Risk Management (NCCR FINRISK). Part of this research was done when he was visiting THEMA and IRES. Downloadable at

http://www.hec.unige.ch/professeurs/SCAILLET\_Olivier/pages\_web/Home\_Page.htm

#### Abstract

In this paper, we characterize explicitly the first derivative of the Value at Risk and the Expected Shortfall with respect to portfolio allocation when netting between positions exists. As a particular case, we examine a simple Gaussian example in order to illustrate the impact of netting agreements in credit risk management. Collateral issues are also dealt with. For practical purposes we further provide nonparametric estimators for sensitivities and derive their asymptotic distributions. An empirical application on a typical banking portfolio is finally provided.

#### Résumé

Dans ce papier, nous calculons les dérivées premières de la "Value at Risk" et de l'"Expected Shortfall", par rapport à de nouvelles positions lorsque ces dernières peuvent faire partie d'un accord de compensation. En particulier, pour illustrer l'impact de la compensation dans un cadre de la gestion du risque, nous examinons le cas simple d'un portefeuille dont les variations de valeurs sont gaussiennes. Nous étendons les formules pour traiter le cas de garanties. Nous fournissons également des estimateurs non paramétriques des sensibilités ainsi que leurs lois asymptotiques. Enfin, une application empirique sur un portefeuille bancaire typique est détaillée.

**Key words:** Value at Risk, Expected Shortfall, Sensitivity, Risk Management, Credit Risk, Netting, Collateral.

JEL Classification: C14, D81, G10, G21, G22.

#### 1 Introduction

For risk management purposes, the evaluation of marginal impacts of current or new positions on risk measures and regulatory capital has been recognized as an important point (Garman (1996), Jorion (1997)). In practice, this evaluation can be made through explicit estimators of the first order derivatives of some standard risk measures, such as the Value at Risk (VaR) and the Expected Shortfall (ES), with respect to portfolio allocations (Gourieroux, Laurent and Scaillet (2000), hereafter GLS, and Scaillet (2000)). Knowledge of the sensitivity is helpful in reducing the amount of computational time needed to process large portfolios since it avoids the need to recompute risk measures each time the portfolio composition is slightly modified (Kurth and Tasche (2002), Martin, Thompson and Brown (2001), Martin and Wilde (2002)). Besides it allows decomposing global portfolio risk component by component, and identifying the largest risk contributions (Denault (2001), Garman (1997), Hallerbach (2003), Tasche (1999)). These derivatives are also of particular relevance in portfolio selection problem (see Markowitz (1952) for portfolio selection in a mean-variance framework). They help to characterize and evaluate efficient portfolio allocations<sup>2</sup> when VaR and ES are substituted for variance as measure of risk (GLS (2000), Rockafellar and Uryasev (2000)). In fact, numerical constrained optimization algorithms for computations of optimal allocations usually require consistent estimates of first order derivatives in order to converge properly.

Unfortunately, the results available up to now have fallen short of takling the problem of netting. Clearly, this is an important omission since most financial positions with respect to one or several counterparties are netted in practice.

Generally speaking, when trading partners agree to offset their positions or obligations, we say that they are netting. By doing so, they reduce a large number of positions or obligations to a smaller number of positions or obligations, and it is on this netted position that the two trading partners settle their outstanding obligations.

In the financial community, positions are most of the time netted inside standardised

<sup>&</sup>lt;sup>2</sup>A related topic is dynamic trading strategies under risk limits (Basak and Shapiro (2001), Cuoco, He, and Issaenko (2001), Leippold, Trojani, and Vanini (2002).

juridical contracts. Streamlining of documentation has taken place as a result of joint efforts by regulators and financial industry organisations. In 1990, the Bank of International Settlements (BIS) issued minimum standards for the design and operation of netting schemes <sup>3</sup>, while in 1991, the Federal Deposit Insurance Corporation Improvement Act (FDICIA) provided support for netting contracts among banks and other financial institutions. In 1992, the International Swaps and Derivatives Dealers (ISDA) issued its first version of the well-known "ISDA Master Agreement" for over-the-counter (OTC) derivatives markets. Its ammended versions are still in force between most market participants around the globe today.

To figure out the relative importance of OTC derivative markets, we recall that the total estimated notional amount of outstanding contracts stood at \$141.7 trillion at the end-of-December 2002 (BIS, 2003), an 11% increase from end-of-June 2002. This compares with a 15% increase in the first half of 2002. At the same time, gross market values grew sharply, rising by 43% to \$6.4 trillion. OTC business continued to accelerate significantly relative to that on exchanges  $^4$ .

As mentioned previously, the term netting is used to describe the process of offsetting mutual positions or obligations e.g. to offset an obligation owed by bank A to bank B with an obligation owed by bank B to bank A. There are three main techniques for netting:

- The payment or settlement netting is the process of settling all deals between two counterparties on a net cash basis, in the same currency. It can be informal or based on a formal agreement. The credit risk between counterparties remains unchanged and they stay legally obligated to settle the gross amounts of their positions. Payment netting allows for reducing the need for intra-day liquidity or credit in bridging the timing gaps between gross payments and gross receipts.
- The netting by novation means that a single net amount is contractually substituted

<sup>&</sup>lt;sup>3</sup>They are known as the Lamfalussy standards after the chairman of the Committee that wrote the report.

<sup>&</sup>lt;sup>4</sup>Open positions on exchange-traded contracts were stagnant, falling by 1% in the second half of the year 2002. It should be noted, however, that activity in the two types of markets cannot be directly and simply compared owing to inherent differences in the characteristics and uses of the products.

for previous individual gross sums owed between two counterparties, i.e. existing obligations are discharged by replacing them with a new obligation. Thus, netting by novation is a formal agreement that aims to reduce liquidity and counterparty credit risk. This novation process may take place automatically within the trading day, on the exchange of confirmations between the two banks. The claims need to be in the same currency. This process can be repeated an infinite number of times until the cut-off time for a particular settlement date. Although netting by novation is, in essence, a bilateral mechanism, it can be operated on multilateral basis within a larger group of banks.

• The close-out netting is an arrangement to settle all contracted but not due liabilities to and claims on an institution by one single payment, immediately after the occurrence of a default or termination event <sup>5</sup>. Since netting by close-out only operates upon the occurrence of a designated event, it cannot have any impact upon the number of payment messages passing between the counterparties in their normal trading relationship. Equally, it has no impact on liquidity or credit risk. If a termination event occurs, each trade is settled individually on due date unless the counterparties also agree on a supplementary netting by novation.

Payment and close-out nettings are part of the ISDA Master agreement. The use of netting techniques can bring significant benefits for balance-sheet purposes, capital usage, credit risk and operational efficiencies <sup>6</sup>. Indeed, it reduces the number of payment messages that have to be exchanged between counterparties. This lowers transaction costs and communication expenses, as well as the chance of mistakes. Moreover, netting is important because it reduces credit and liquidity risks, and ultimately systemic risk. Taking an offsetting position subject to a netting agreement is thus related to credit risk mitigation. The tendency of the regulator is to allow wider range of credit risk mitigants in order to avoid the so-called "domino effect" in the financial sector. In the new Basel Capital

<sup>&</sup>lt;sup>5</sup>An early termination date is designated as the result of the occurrence of an event of default or a decline in the credit quality of a party following the occurrence of merger-related events.

<sup>&</sup>lt;sup>6</sup>See also DUFFIE and SINGLETON (2003) for impact of netting in credit derivatives asset pricing.

Accord (see the 2001 consultative document), on-balance sheet netting agreements of loans and deposits of banks to or from a counterparty will be permitted under some conditions. Note that, in 1995, the 1988 Basel Accord was modified to allow banks to reduce the credit exposures ("credit equivalent" in the Basel terminology) of their derivative positions by bilateral netting procedures (see Crouhy et al. (1998)).

The paper is organised as follows. In Sections 2 and 3 our aim is to extend the sensitivity analysis of VaR and ES to a setting in which netting is allowed, and to propose suitable estimators of the first order derivatives of VaR and ES in that context. In Section 2, we outline the framework and explain the differences arising from netting agreements in case of default of a given counterparty. The loss function associated with netted positions is no more a simple sum of exposures or mark-to-market valuations, but rather involves some nonlinearities. More precisely, it involves some terms like  $(Y_1 + \ldots + Y_I)^+ = \max(Y_1 + \ldots + Y_I)^+$  $Y_{I}, 0$ , when the positions  $Y_{i}, i = 1, ..., I$ , belong to the same netting agreement at default. Section 2 contains the main result of the paper, namely the explicit characterization of the first order derivatives of VaR and ES for portfolios under netting agreements. The Gaussian case is briefly discussed. In particular we compare VaR and its sensitivity in the netted and unnetted cases, and show on a simple stylized example that netting is a valuable credit risk mitigation device. In Section 3 we outline an extension of the approach, which allows for the presence of collateral. In Section 4, we derive estimators of sensitivities so that they can be used in practical risk management and portfolio selection procedures. These estimators are of a nonparametric nature and easy to implement. In Section 5 we provide an empirical illustration for a typical portfolio of a large bank. Section 6 contains some concluding remarks. Technical appendices gather proofs.

#### 2 Main results

Let us consider a portfolio made of three components: a'Y,  $a'_UY_U$  and  $a'_NY_N$ . The first component a'Y is the part which is not sensitive to default risk on a given counterparty, while the two other parts can be affected by its default <sup>7</sup>. The second component  $a'_U Y_U$ gathers unnetted positions, i.e. positions which are outside the netting agreement signed with the counterparty and will not be pooled into a single position in case of counterparty default. The third component  $a'_N Y_N$  is the netted part governed by the master agreement with the given counterparty. At default, unnetted and netted positions should receive a different treatment when they exhibit positive value. Indeed recall that only contracts showing positive value at default need to be included in credit loss computations. The credit loss on unnetted positions corresponds to the sum on the positive part of each position, i.e.  $\sum_j (a_{j,U}Y_{j,U})^+$ , while the credit loss on netted positions corresponds to the positive part of the sum on each position, i.e.  $(a'_N Y_N)^+$ .

Let us now consider an additional single position  $a_Z Z$ . This leads to a portfolio value at the initial date  $t_0$  equal to:

$$V(t_0) = a'Y(t_0) + a'_U Y_U(t_0) + a'_N Y_N(t_0) + a_Z Z(t_0).$$

Hereafter the random variable D indicates default of the given counterparty at date  $t_1$ , and takes the value 1 (resp. 0) in case of default (resp. no default). More generally, Dcorresponds to a credit event that causes some losses, and is the starter of a netting agreement.

In the derivation of the portfolio value at date  $t_1$  we need to distinguish three cases: i) the position is not subject to default risk on the counterparty; ii) the position is subject to default risk and cannot be netted in case of counterparty default; iii) the position is subject to default risk and falls under the umbrella of the master agreement signed with the counterparty. Then the values of the global portfolio at  $t_1$  are obtained from the contributions of each component under default and no default after taking into account potential netting <sup>8</sup>.

<sup>&</sup>lt;sup>7</sup>Note that the identification of default to a given single counterparty is not a limitation of our analysis. The setting is sufficiently large to accommodate the practical situation of portfolios subject to multiple defaults.

<sup>&</sup>lt;sup>8</sup>In case of default it may happen that the asset Z urgently needs to be replaced because of hedging demand. It is then sufficient to substract the replacement value in case of default at date  $t_1$ , i.e. add the short position  $-a_Z Z(t_1)D$ . Stated results may be easily adapted to handle that case.

i) No default risk on  $a_Z Z$ 

$$V(t_1) = a'Y(t_1) + a_Z Z(t_1) + (a'_U Y_U(t_1) + a'_N Y_N(t_1)) - \left(\sum_j (a_{j,U} Y_{j,U}(t_1))^+ + (a'_N Y_N(t_1))^+\right) D.$$

ii) Default risk on unnetted  $a_Z Z$ 

$$V(t_1) = a'Y(t_1) + (a'_U Y_U(t_1) + a_Z Z(t_1) + a'_N Y_N(t_1)) - \left(\sum_j (a_{j,U} Y_{j,U}(t_1))^+ + (a_Z Z(t_1))^+ + (a'_N Y_N(t_1))^+\right) D.$$

iii) Default risk on netted  $a_Z Z$ 

$$V(t_1) = a'Y(t_1) + (a'_UY_U(t_1) + a'_NY_N(t_1) + a_ZZ(t_1)) - \left(\sum_j (a_{j,U}Y_{j,U}(t_1))^+ + (a'_NY_N(t_1) + a_ZZ(t_1))^+\right) D.$$

In the following, we are interested in risk measures computed on the portfolio value at date  $t_1$ , namely VaR and ES (see Szegö (2002) for a discussion about conditions for proper use of risk measures) defined by:

$$\alpha = P[V(t_1) < -VaR], \tag{1}$$

$$ES = -E[V(t_1)|V(t_1) < -VaR],$$
(2)

where  $\alpha$  is a small loss probability level, say 1% <sup>9</sup>. The dependence of VaR and ES with respect to the loss probability level and portfolio allocations will be assumed implicit in the notations. Equations (1) and (2) may also be expressed in terms of losses using L = -V:

$$\alpha = P[L(t_1) > VaR], \tag{3}$$

$$ES = E[L(t_1)|L(t_1) > VaR].$$
 (4)

<sup>9</sup>These risk measures may also be computed on portfolio value *changes*:

$$\alpha = P[V(t_1) - V(t_0) < -\overline{V}aR],$$
  
$$\overline{E}S = -E[V(t_1) - V(t_0)|V(t_1) - V(t_0) < -\overline{V}aR].$$

Results obtained for VaR and ES can be immediately translated to  $\overline{V}aR$  and  $\overline{E}S$  using the two changes of variable  $\overline{V}aR = VaR + V(t_0)$  and  $\overline{E}S = ES + V(t_0)$ . In particular, this will lead to  $\frac{\partial \overline{V}aR}{\partial a_Z} = \frac{\partial VaR}{\partial a_Z} + \frac{\partial V(t_0)}{\partial a_Z}$  and  $\frac{\partial \overline{E}S}{\partial a_Z} = \frac{\partial ES}{\partial a_Z} + \frac{\partial V(t_0)}{\partial a_Z} = Z(t_0)$ .

Note that our definition of "Expected Shortfall" does not correspond to the most general definition of this coherent measure of risk (see e.g. Acerbi and Tasche (2002), Rockafellar and Uryasev (2002)):

$$\tilde{ES} = E[L(t_1)|L(t_1) > VaR] - (\alpha - P(L(t_1) > VaR))\frac{1}{\alpha}VaR.$$

Nonetheless, under our assumptions, the law of the loss functions will be absolutely continuous with respect to the Lebesgue measure. In this case, we know that ES and  $\tilde{ES}$ coincide.

As already mentioned, it is interesting to compute the contribution of each position to the risk of the whole portfolio and monitor the most risky ones for precautionary reasons. Since VaR and ES are homogeneous of degree one, we get by Euler Theorem (see Tasche (1999), for instance) :

$$a'\frac{\partial VaR}{\partial a'} + a_Z \frac{\partial VaR}{\partial a'_Z} + a'_U \frac{\partial VaR}{\partial a'_U} + a'_N \frac{\partial VaR}{\partial a'_N} = VaR,$$

and the same expression holds with ES sustituted for VaR. The contribution associated with  $a_Z Z$  to the global risk measured by VaR (resp. ES) is simply the sensitivity of VaR (resp. ES) with respect to  $a_Z$  multiplied by the allocation  $a_Z$ , i.e.  $a_Z \partial VaR/\partial a_Z$ , resp.  $a_Z \partial ES/\partial a_Z$ . Such a quantity is known in the literature (see e.g. Garman (1997)) as incremental or component VaR.

The first order derivative of VaR and ES with respect to  $a_Z$  can be easily computed by applying the following proposition.

**Proposition 1.** Consider the loss functions

$$L_1 = X - \varepsilon Z,$$
  

$$L_2 = X - \varepsilon Z + (\varepsilon Z)^+ D,$$
  

$$L_3 = X - \varepsilon Z + (Y + \varepsilon Z)^+ D,$$

where  $\varepsilon$  is a positive real number <sup>10</sup>, and (X, Y, Z) is a random vector admitting a continuous conditional density with respect to the Lebesgue measure, conditionally to the event

<sup>&</sup>lt;sup>10</sup> if  $\varepsilon$  is negative, the results can be applied by simply changing Z into -Z.

D = 1 and D = 0. Then

A. First derivative of Value at Risk.

$$\partial_{\varepsilon} VaR_1 = -E[Z|L_1 = VaR_1], \tag{5}$$

$$\partial_{\varepsilon} VaR_2 = -E[Z\mathbf{1}\{Z<0\}|L_2 = VaR_2, D = 1] \times P[D = 1|L_2 = VaR_2] -E[Z|L_2 = VaR_2, D = 0] \times P[D = 0|L_2 = VaR_2],$$
(6)

$$\partial_{\varepsilon} VaR_3 = -E[Z\mathbf{1}\{Y + \varepsilon Z < 0\} | L_3 = VaR_3, D = 1] \times P[D = 1 | L_3 = VaR_3] -E[Z|L_3 = VaR_3, D = 0] \times P[D = 0 | L_3 = VaR_3].$$
(7)

#### B. First derivative of Expected Shortfall.

$$\partial_{\varepsilon} ES_{1} = -E[Z|L_{1} > VaR_{1}], \tag{8}$$

$$\partial_{\varepsilon} ES_{2} = -E[Z\mathbf{1}\{Z < 0\}|L_{2} > VaR_{2}, D = 1] \times P[D = 1|L_{2} > VaR_{2}] -E[Z|L_{2} > VaR_{2}, D = 0] \times P[D = 0|L_{2} > VaR_{2}], \tag{9}$$

$$\partial_{\varepsilon} ES_3 = -E[Z\mathbf{1}\{Y + \varepsilon Z < 0\} | L_3 > VaR_3, D = 1] \times P[D = 1 | L_3 > VaR_3] -E[Z|L_3 > VaR_3, D = 0] \times P[D = 0 | L_3 > VaR_3].$$
(10)

Remark 1. Note that we can rewrite simply

$$\begin{array}{lll} \partial_{\varepsilon} VaR_{2} &=& -E[Z\mathbf{1}\{Z<0,D=1\}|L_{2}=VaR_{2}]-E[Z\mathbf{1}\{D=0\}\}|L_{2}=VaR_{2}],\\ \partial_{\varepsilon} VaR_{3} &=& -E[Z\mathbf{1}\{Y+\varepsilon Z<0,D=1\}|L_{3}=VaR_{3}]-E[Z\mathbf{1}\{D=0\}\}|L_{3}=VaR_{3}], \end{array}$$

and

$$\begin{array}{lll} \partial_{\varepsilon} ES_2 &=& -E[Z\mathbf{1}\{Z<0,D=1\}|L_2>VaR_2]-E[Z\mathbf{1}\{D=0\}|L_2>VaR_2],\\ \partial_{\varepsilon} ES_3 &=& -E[Z\mathbf{1}\{Y+\varepsilon Z<0,D=1\}|L_3>VaR_3]-E[Z\mathbf{1}\{D=0\}|L_3>VaR_3]. \end{array}$$

In the aforementioned loss functions we have implicitly assumed zero recovery rates. This point of view, not necessarily unrealistic, could appear as too conservative in some cases. Similar formulas to the ones presented in Proposition 1 may be obtained with arbitrary fixed or random recovery rates. Unfortunately, these formulas are a lot more complicated. Indeed, with a non zero recovery rate R, the loss function, say  $L_3$ , becomes  $X - \varepsilon Z + (1 - R)(Y + \varepsilon Z)^+$ . When differentiating with respect to  $\varepsilon$ , the simplification between the terms that are coming from the market value  $X - \varepsilon Z$  and those that are coming from the credit exposure  $(1 - R)(Y + \varepsilon Z)^+$  disappears. Fortunately this problem is avoided when dealing with credit exposures only, as we will see below.

Previous computations have been made in an "integrated" credit and market risk environment. Indeed, Var and ES computations have included impact of both market conditions and credit events on the portfolio value. The previous random variables X, Y, Zare usually functions of some market factors (equity indices, interest rates, exchange rates, credit spreads, ...). Let us remark that we do not rely on the assumption of independence between credit events and changes in these underlying market factors when deriving our results. Actually, some standard credit risk models are based on a dependence between credit events and the variations of some underlying market factors such as equity prices ("structural approaches") or spreads.

For credit analysis purposes, it is common to focus on credit risk solely. Practitioners and regulators are then concerned with so-called exposures to default. The exposure is a random variable related to positions with respect to a single counterparty, say i, and corresponds to a specific amount potentially lost due to the default of that counterparty in the future. Lost amounts are often calculated by some calibrated pricing models. Formally, the exposure at  $t_1$  to counterparty i can be written as

$$Exp_{U,i}(t_1) = \left(\sum_{j} (a_{j,U}Y_{j,U}(t_1))^+ + (a_Z Z(t_1))^+ + (a'_N Y_N(t_1))^+\right), \quad (11)$$

in the unnetted case, and

$$Exp_{N,i}(t_1) = \left(\sum_{j} (a_{j,U}Y_{j,U}(t_1))^+ + (a'_NY_N(t_1) + a_ZZ(t_1))^+\right),$$
(12)

in the netted case.

Let us stress that positions are related to the same single counterparty i. In the first case, Z is outside a netting agreement while, in the second case, it is not. Note further that exposures have nothing to do with default probabilities of the counterparty. The VaR associated with a single exposure is commonly called a potential exposure, and is a pure credit risk measure linked to positions with a single counterparty.

Moreover, to compute provisions to cover multiple credit losses, it is necessary to aggregate exposures and take into account the probability of default of several counterparties. Losses can be aggregated, for example, inside some portfolios, sections, or perimeters. The total credit loss of a portfolio  $\mathcal{P}$  at  $t_1$  is then equal to

$$L(t_1) = \sum_{i \in \mathcal{P}} Exp_i(t_1)D_i,$$

denoting by  $D_i$  the default of the *i*-th counterparty at date  $t_1$ . To study the marginal impact of a particular position Z related to a given counterparty *i* on the VaR and ES of the portfolio  $\mathcal{P}$  is approximately equivalent, depending on the status of Z in terms of netting, to calculate the sensitivity to  $a_Z$  of the VaR and ES of

$$L_U(t_1) = X + \left(\sum_j (a_{j,U}Y_{j,U}(t_1))^+ + (a_Z Z(t_1))^+ + (a'_N Y_N(t_1))^+\right) D_i,$$

and

$$L_N(t_1) = X + \left(\sum_j (a_{j,U}Y_{j,U}(t_1))^+ + (a'_NY_N(t_1) + a_ZZ(t_1))^+\right) D_i,$$

for some random variable X.

These sensitivities can be calculated using the following result.

**Proposition 2.** Consider the loss functions

$$L_4 = X + (\varepsilon Z)^+ D,$$
  

$$L_5 = X + (Y + \varepsilon Z)^+ D,$$

where  $\varepsilon$  is a positive real number, and (X, Y, Z) is a random vector admitting a continuous conditional density with respect to the Lebesgue measure, conditionally to the event D = 1 and D = 0. Then

A. First derivative of Value at Risk.

$$\partial_{\varepsilon} VaR_4 = E[Z\mathbf{1}\{Z \ge 0\} | L_4 = VaR_4, D = 1] \times P[D = 1 | L_4 = VaR_4],$$
(13)

$$\partial_{\varepsilon} VaR_5 = E[Z\mathbf{1}\{Y + \varepsilon Z \ge 0\}|L_5 = VaR_5, D = 1] \times P[D = 1|L_5 = VaR_5].$$
 (14)

B. First derivative of Expected Shortfall.

$$\partial_{\varepsilon} ES_4 = E[Z\mathbf{1}\{Z \ge 0\} | L_4 > VaR_4, D = 1]P[D = 1 | L_4 > VaR_4],$$
(15)

$$\partial_{\varepsilon} ES_5 = E[Z\mathbf{1}\{Y + \varepsilon Z > 0\} | L_5 > VaR_5, D = 1]P[D = 1 | L_5 > VaR_5].$$
(16)

**Remark 2.** Note that we can rewrite

$$\partial_{\varepsilon} VaR_4 = E[Z\mathbf{1}\{Z \ge 0, D=1\}|L_4 = VaR_4],$$
  
$$\partial_{\varepsilon} VaR_5 = E[Z\mathbf{1}\{Y + \varepsilon Z \ge 0, D=1\}|L_5 = VaR_5],$$

and

$$\begin{aligned} \partial_{\varepsilon} ES_4 &= E[Z\mathbf{1}\{Z > 0, D = 1\} | L_4 > VaR_4], \\ \partial_{\varepsilon} ES_5 &= E[Z\mathbf{1}\{Y + \varepsilon Z > 0, D = 1\} | L_5 > VaR_5]. \end{aligned}$$

Clearly sensitivities of potential exposures as defined in Equations (11) and (12) can be deduced from the previous formulas by forcing D = 1. The above formulas can also be easily modified to incorporate the presence of a non zero (constant or random) recovery rate.

All aforementioned expressions are explicit. In principle they can be computed, at least numerically, if the joint distribution of (X, Y, Z, D) is known. The exposures are clearly functions of some underlying market factors. Thus, as we have said previously, (X, Y, Z)may depend on D. Nonetheless, once the distribution of the underlying random variables (X, Y, Z) is known and independence with respect to the default event is assumed, estimation is much more simple. This will be the case for reduced form credit risk models where default is not predictable and whose law does not depend on the current or past values of market factors. Implicitly, we will do this assumption in our numerical illustration.

For instance, assume the joint law of (X, Y, Z) is Gaussian  $\mathcal{N}(m, \Omega)$ . Then it possible to get a closed form expression for  $\partial_{\varepsilon} VaR_j$  and  $\partial_{\varepsilon} ES_j$ ,  $j = 1, \ldots, 5$ . Nevertheless, the formulas are tedious to obtain since the positive parts in the losses functions  $L_j$  require to evaluate some truncated distributions.

The assumption of normality may seem odd to some readers: the exposures are usually sums of more or less nonlinear instruments with respect to market factors, and these factors themselves often exhibit non Gaussian features (fat tails, volatility clustering). Normality can be understood under a standard market risk approach, where exposures are sums of hundreds of net values. Alternatively, when the time horizon is growing, the distributions of exposures tend to be Gaussian. Thus, we could consider market exposures in the long run, i.e. for a time horizon of several years, to justify this type of assumption.

In the remaining of this section, we only study a simple portfolio example  $E_1 + \epsilon E_2$ where the vector of exposures  $E = (E_1, E_2)'$  is Gaussian with mean  $m = (m_1, m_2)'$  and covariance matrix

$$\Omega = \begin{bmatrix} \sigma_1^2 & \rho \sigma_1 \sigma_2 \\ \rho \sigma_1 \sigma_2 & \sigma_2^2 \end{bmatrix}.$$

This stylized example is used to illustrate the impact of netting on VaR and their sensitivities.

Let us consider the credit losses when there is no netting agreement between  $E_1$  and  $E_2$ :

$$L_U = (E_1^+ + (\epsilon E_2)^+)D,$$

and when all positions are netted:

$$L_N = (E_1 + \epsilon E_2)^+ D.$$

The credit VaR themselves do not admit explicit forms (even in this simple case) and need to be computed numerically. However, once they are known, they can be plugged in the explicit form of the sensitivities derived in appendix D.

Figure 1 shows the impact of netting as a function of  $\rho$  for  $m_1 = 10$ ,  $m_2 = -1$ ,  $\varepsilon = 1$ and  $\sigma_1 = \sigma_2 = 1$ . This portfolio can be viewed as made of a long aggregated position in one category of assets and a short position in another one with the same counterparty. In the first case the two positions cannot be netted while they can in the second case. VaR are estimated by computing empirical quantiles on 100,000 simulations delivered by an antithetic variate Monte Carlo procedure. The loss probability level  $\alpha$  is equal to 1% (99% VaR) and the default probability p is equal to 20%.

- Please insert Figure 1 approximately here -

It can be observed that netting is valuable from a credit risk management point of view. Indeed, VaR are always lower in the netted case <sup>11</sup>. These results show that netting can be thought as sound credit risk mitigation practice and should be encouraged. Moreover, the range of VaR sensitivities is larger in the netted case. Hence this example also shows that the sensitivities can differ substantially between the two situations, and that netting agreements should be taken into account when computing VaR sensitivities. In the netted case we may observe that the VaR sensitivity becomes negative for highly negative correlations. The explanation is simple. If  $E_1$  and  $E_2$  are strongly negatively correlated,  $E_2$  is usually very negative when  $E_1$  is very positive. Hence increasing slightly the position in  $E_2$  has a favourable effect on the VaR of the global netted position. On the contrary it is not possible to benefit from the offsetting effect in the unnetted case, and the VaR sensitivity stays positive always.

#### **3** Extension to collateral agreements

Netting agreements are only one part of the credit mitigation techniques. The possibility to exchange collateral is often used by lots of dealers. It is recognized as a way to reduce significantly the amount of credit exposures, even if it may entail other risks: liquidity risk, legal risk, custody risk and operational risk.

The ISDA credit forms - each referred to as a Credit Support Annex (CSA) - that go with international master agreements are now part of the standard documentation. Most national master agreements also have a specific collateral annex. The CSA includes a schedule of elections to be negociated between the counterparties, including a counterparty threshold, that is the amount of uncollateralized credit exposure the other is willing to accept. In general, the threshold reflects the counterparty credit rating: the higher the rating, the larger the threshold.

The collateral reduces the current exposure of the collateral taker to the collateral giver by the amount of collateral held. Even if the net value of the collateral is varying, we

<sup>&</sup>lt;sup>11</sup>Note that if we increase the mean of  $E_2$  in absolute terms, the benefits of netting increase since the probability of taking positive values diminishes.

will assume for the sake of computational convenience that the collateral process bounds the net exposure from above by some constant threshold M. Thus, the loss functions  $L_j, j = 1, \ldots, 5$  stay unchanged except for the positivity condition being replaced by the belonging to the interval [0, M]. The new sensitivities can be calculated as previously. With obvious notations we provide the results for the most general loss functions only.

**Proposition 3.** Consider the loss functions

$$\begin{split} L'_3 &= X - \varepsilon Z + (Y + \varepsilon Z). \mathbf{1} \{ Y + \varepsilon Z \in [0, M] \} D + M. \mathbf{1} \{ Y + \varepsilon Z > M \} D, \\ L'_5 &= X + (Y + \varepsilon Z). \mathbf{1} \{ Y + \varepsilon Z \in [0, M] \} D + M. \mathbf{1} \{ Y + \varepsilon Z > M \} D, \end{split}$$

where  $\varepsilon$  is a positive real number, and (X, Y, Z) is a random vector admitting a continuous conditional density with respect to the Lebesgue measure, conditionally to the event D = 1 and D = 0. Then

A. First derivative of Value at Risk.

$$\partial_{\varepsilon} VaR'_{3} = -E[Z\mathbf{1}\{Y + \varepsilon Z \notin [0, M], D = 1\} | L'_{3} = VaR'_{3}] \\ -E[Z\mathbf{1}\{D = 0\} | L'_{3} = VaR'_{3}],$$
(17)

$$\partial_{\varepsilon} VaR'_5 = E[Z\mathbf{1}\{Y + \varepsilon Z \in [0, M], D = 1\}|L'_5 = VaR'_5].$$

$$(18)$$

#### B. First derivative of Expected Shortfall.

$$\partial_{\varepsilon} ES'_{3} = -E[Z\mathbf{1}\{Y + \varepsilon Z \notin [0, M], D = 1\} | L'_{3} > VaR'_{3}] \\ -E[Z\mathbf{1}\{D = 0\} | L'_{3} > VaR'_{3}],$$
(19)

$$\partial_{\varepsilon} ES'_5 = E[Z\mathbf{1}\{Y + \varepsilon Z \in [0, M], D = 1\} | L'_5 > VaR'_5].$$

$$(20)$$

Note that the formulas are remarkably similar to previous ones. This was not obvious from the start, in particular for  $L'_3$ .

#### 4 Estimation

Since all sensitivities can be evaluated through conditional expectations, they are amenable to estimation by nonparameric techniques such as the well-known kernel method. To be specific, assume the joint distribution of (X, Y, Z, D) is known explicitly, or that we can get at least *n* random draws  $(x_i, y_i, z_i, d_i)_{i=1,...,n}$  from the joint distribution of (X, Y, Z, D). This is the usual situation because most of the risk models are parametric and/or simulation-based. This setting is large enough to host some dependence between "the market factors" (X, Y, Z) and the default event <sup>12</sup>.

Let us consider a consistent estimator  $\widehat{VaR_4}$  of  $VaR_4$ . For instance, it can be deduced from computing an empirical quantile on the simulated losses  $l_{4,i}$ , i = 1, ..., n. Then, an estimate of  $\partial_{\varepsilon} VaR_4$  can be defined by the usual nonparametric kernel regression estimator:

$$\widehat{\partial_{\varepsilon} VaR_4} = \frac{\sum_{i=1}^n z_i \mathbf{1}\{z_i \ge 0, d_i = 1\} K_h(l_{4,i} - \widehat{VaR_4})}{\sum_{i=1}^n K_h(l_{4,i} - \widehat{VaR_4})},$$
(21)

where  $K : \mathbb{R} \to \mathbb{R}$  is a kernel (an integrable function whose integral is one), h = h(n) is a smoothing parameter, called the bandwidth (a sequence of positive real numbers which tends to 0 when n tends to the infinity) and  $K_h(u) = K(u/h)/h^{-13}$ . Similarly,

$$\widehat{\partial_{\varepsilon} VaR_5} = \frac{\sum_{i=1}^n z_i \mathbf{1}\{y_i + \varepsilon z_i \ge 0, d_i = 1\} K_h(l_{5,i} - \widehat{VaR_5})}{\sum_{i=1}^n K_h(l_{5,i} - \widehat{VaR_5})}.$$
(22)

Both estimators are easy to implement since they correspond to ratios of straight averages of known functions of the observed data points.

Denote the density of  $L_k$  by  $f_k$ , k = 4, 5, and set

$$\mu_4(v) = E[Z^2 \mathbf{1}\{Z > 0, D = 1\} | L_4 = v] - (\partial_{\varepsilon} V a R_4)^2,$$
  
$$\mu_5(v) = E[Z^2 \mathbf{1}\{Y + \varepsilon Z > 0, D = 1\} | L_5 = v] - (\partial_{\varepsilon} V a R_5)^2.$$

Then we prove the following asymptotic normality result in Appendix D.

**Proposition 4.** Assume for k = 4, 5,

- $\sqrt{n}(\widehat{VaR_k} VaR_k) = O_P(1),$
- K is an even kernel function,  $\int |t|^3 |K|(t) dt < \infty$ ,  $\lim_{|t|\to\infty} |t|^3 K(t) = 0$ . It is three times continuously differentiable, K' and K'' are integrable and K''' is bounded.

 $<sup>^{12}\</sup>mathrm{For}$  instance this dependence exists when equity prices and default probabilities are linked.

<sup>&</sup>lt;sup>13</sup>Obviously, direct differentiation of  $\widehat{VaR_4}$  through computation of finite difference has no meaning here since the empirical quantile is *not* differentiable w.r.t.  $\epsilon$ .

- $E[|Z|^p] < \infty$  for every integer p,
- $t \mapsto E[Z^4|L_k = t]$  is bounded in a neighborhood of  $VaR_k$ ,
- $nh^5 \longrightarrow 0$  and  $nh^{7/2} \longrightarrow \infty$ ,
- $f_k$  and the function  $VaR_k \mapsto \partial_{\varepsilon} VaR_k$  are two times continuously differentiable,
- $f_k$  and  $\mu_k$  are continuous and strictly positive in a neighborhood of  $VaR_k$ .

Then

$$\begin{split} \sqrt{nh} \left\{ \widehat{\partial_{\varepsilon} VaR_k} - \partial_{\varepsilon} VaR_k \right\} \xrightarrow[n \to \infty]{law} \mathcal{N}(0, \sigma_k^2). \\ \sigma_k^2 &= \frac{\int K^2 \cdot \mu_k(VaR_k)}{f_k(VaR_k)}. \end{split}$$

Similar kernel estimators of  $\partial_{\varepsilon} VaR_j$ , j = 1, 2, 3 can be easily inferred. They are asymptotically normal under similar regularity conditions.

As usually these asymptotic normality results can be used to build confidence intervals around estimates. In particular the number n of simulations can be chosen so that these intervals are sufficiently narrow to guarantee statistically precise sensitivity estimates. The previous asymptotic variances  $\sigma_k^2$  can be easily estimated. For instance, to get an estimate of  $\sigma_5^2$ , we may replace  $\mu_5(VaR_5)$  by

$$\hat{\mu}_{5} = \frac{\sum_{i=1}^{n} z_{i}^{2} \mathbf{1} \{ y_{i} + \varepsilon z_{i} \ge 0, d_{i} = 1 \} K_{h} (l_{5,i} - \widehat{VaR_{5}})}{\sum_{i=1}^{n} K_{h} (l_{5,i} - \widehat{VaR_{5}})} - \left( \widehat{\partial_{\varepsilon} VaR_{5}} \right)^{2},$$

and  $f_5(VaR_5)$  by

$$\hat{f}_5 = \frac{1}{n} \sum_{i=1}^n K_h (l_{5,i} - \widehat{VaR_5}).$$

We can also estimate the sensitivity of the expected shortfall by

$$\widehat{\partial_{\varepsilon} ES_4} = \frac{\sum_{i=1}^n z_i \mathbf{1}\{z_i \ge 0, d_i = 1, l_{4,i} > \widehat{VaR_4}\}}{\sum_{i=1}^n \mathbf{1}\{l_{4,i} > \widehat{VaR_4}\}},$$
(23)

and

$$\widehat{\partial_{\varepsilon} ES_5} = \frac{\sum_{i=1}^n z_i \mathbf{1}\{y_i + \varepsilon z_i \ge 0, d_i = 1, l_{5,i} > \widehat{VaR_5}\}}{\sum_{i=1}^n \mathbf{1}\{l_{5,i} > \widehat{VaR_5}\}}.$$
(24)

The strong consistency of the two latter estimators can be deduced easily from the strong uniform consistency of the empirical process (see Van der Vaart (1994)). Furthermore, their asymptotic normality can be established under some regularity conditions. However the forms of their asymptotic normal distributions depend on the choice of the estimators  $\widehat{VaR_k}$ , and are more complicated than in the VaR case.

Clearly, the previous estimation procedures are straighforward to modify to account for the presence of collateral. We only have to replace the indicator functions  $\mathbf{1}\{y_i + \varepsilon z_i > 0\}$ by  $\mathbf{1}\{y_i + \varepsilon z_i \in [0, M]\}$  as before.

#### 5 Empirical application

In this empirical section, we wish to illustrate how the estimators of the previous section can be used in practice. We study a real life banking example and analyze the sensitivity w.r.t. an additional short position in a bond (borrowing) with a single AA- counterparty. The default probability is 2% at one year. We examine a short position to emphasize importance of netting in credit risk calculations, and aim to quantify what is the impact of this additional position on credit VaR and credit ES at a 1% loss probability level. 35 other positions are already standing with this counterparty. Each of these 35 positions corresponds to an aggregated position inside a netting agreement (some of them are in fact stand alone positions). Positions can be long or short, and include several categories of assets such as forward contracts, swaps and options in different currencies. Table 1 gives the current value of the 35 aggregated positions in Euros. The current value of the additional short position is equal to -1,150,000 Euros.

- Please insert Table 1 approximately here -

We compare the situation where the additional position can be incorporated inside one of the 35 existing netting agreements with the case where it is left outside any netting <sup>14</sup>. The 35 netted cases are numbered from 1 to 35, and the unnetted case is called case 0.

 $<sup>^{14}</sup>$ Let us remark that this illustrative example may look a bit fictitious since one cannot usually choose the status of an additional position. We do so because this enlarges the number of studied cases.

The number of simulations of default times is n = 100,000, and random future position values are computed from internal pricing models. The estimation procedure is based on a Gaussian kernel and a bandwidth selected according to the usual rule of thumb, namely  $h = \hat{\sigma} n^{-1/5}$  where  $\hat{\sigma}$  is the empirical standard deviation of the credit losses. Table 2 gives the one year credit VaR and one year credit ES together with their sensitivity for each case.

#### - Please insert Table 2 approximately here -

As already observed on the Gaussian numerical example, credit VaR and credit ES may differ substantially between netted and unnetted cases. Credit ES are larger than credit VaR revealing much of the danger of using the VaR concept in a credit environment (see Frey and McNeil (2002) for further elaboration). Besides if one enjoys the possibility to choose the agreement, which will host the additional position, then cases 17, 26, 27, 30, 35, resp. 28, seem to be more suitable from a credit risk management point of view since they minimize VaR, resp. ES, at one year. It can be further observed that a small VaR sensitivity can coexist with a larger ES sensitivity (see for instance case 15). This can be explained by the fact that VaR only concerns a single point in the loss distribution, while ES is about an average of points.

#### 6 Concluding remarks

Risk measures answer the need of quantifying the risk of potential losses on a portfolio of assets. This need may arise due to internal concerns (risk-reward tradeoff) or external constraints (prudential rules imposed by regulators). In this paper we have proposed estimation procedures allowing for a sensivity analysis w.r.t. changes in the portfolio allocation. The setting explicitly takes into account the possibility of netting and collateral agreements. The estimation procedures are nonparametric, fast and easy to implement. They have also been shown to be of practical relevance in real life banking situations, and should help to achieve better credit risk management in the future.

1	2	3	4	5
9,100,000	510	$752,\!076$	556,260	-2,140,000
6	7	8	9	10
$-2.510 \ 10^9$	-17,100,000	-22,500,000	3,990,000	-1,520,000
11	12	13	14	15
654,270	$688,\!179$	6,890,000	893,581	-28,000,000
16	17	18	19	20
-1,190,000	30,100,000	$156,\!448$	$-1,\!680,\!000$	0
21	22	23	24	25
0	0	-29,400,000	-2,010,000	3,220,000
26	27	28	29	30
-872,964	24,900,000	3,700,000	388,944	3,850,000
31	32	33	34	35
-346,792	294,316	-77,199	-1,620,000	3,340,000

Table 1: Current aggregated position values

case	VaR	$\widehat{\partial_{\varepsilon} VaR}$	$\widehat{ES}$	$\widehat{\partial_{\varepsilon} ES}$
0	$7.033 \ 10^7$	0	$8.380 \ 10^7$	0
1	$6.937 \ 10^7$	$-0.087 \ 10^7$	$8.304  10^7$	$-0.093 \ 10^7$
2	$7.033 \ 10^7$	0	$8.379  10^7$	0
3	$6.954 \ 10^7$	$-0.062 \ 10^7$	$8.318 \ 10^7$	$-0.060 \ 10^7$
4	$7.007 \ 10^7$	$-0.003 \ 10^7$	$8.371  10^7$	$-0.002 \ 10^7$
5	$7.033 \ 10^7$	0	$8.380 \ 10^7$	0
6	$7.033 \ 10^7$	0	$8.380 \ 10^7$	0
7	$7.033 \ 10^7$	0	$8.380 \ 10^7$	0
8	$7.033 \ 10^7$	0	$8.380 \ 10^7$	0
9	$6.981 \ 10^7$	0	$8.329 \ 10^7$	0
10	$7.001 \ 10^7$	$-0.015 \ 10^7$	$8.354 \ 10^7$	$-0.023 \ 10^7$
11	$7.019 \ 10^7$	0	$8.365 \ 10^7$	0
12	$7.019 \ 10^7$	$-0.005 \ 10^7$	$8.346 \ 10^7$	$-0.013 \ 10^7$
13	$6.941 \ 10^7$	$-0.069 \ 10^7$	$8.301  10^7$	$-0.083 \ 10^7$
14	$7.001 \ 10^7$	$-0.008 \ 10^7$	$8.350 \ 10^7$	$-0.018 \ 10^7$
15	$7.033 \ 10^7$	$-0.004 \ 10^7$	$8.359 \ 10^7$	$-0.019 \ 10^7$
16	$6.975 \ 10^7$	0	$8.332 \ 10^7$	0
17	$6.928 \ 10^7$	$-0.101 \ 10^7$	$8.283 \ 10^7$	$-0.103 \ 10^7$
18	$7.017 \ 10^7$	0	$8.364  10^7$	0
19	$7.033 \ 10^7$	0	$8.380 \ 10^7$	0
20	$7.033 \ 10^7$	0	$8.380 \ 10^7$	0
21	$7.033 \ 10^7$	0	$8.380 \ 10^7$	0
22	$7.033 \ 10^7$	0	$8.380 \ 10^7$	0
23	$7.029 \ 10^7$	$-0.002 \ 10^7$	$8.382 \ 10^7$	$-0.003 \ 10^7$
24	$6.962 \ 10^7$	$-0.033 \ 10^7$	$8.328 \ 10^7$	$-0.038 \ 10^7$
25	$7.033 \ 10^7$	0	$8.380 \ 10^7$	0
26	$6.928 \ 10^7$	$-0.080 \ 10^7$	$8.291  10^7$	$-0.087 \ 10^7$
27	$6.928 \ 10^7$	$-0.101 \ 10^7$	$8.283 \ 10^7$	$-0.103 \ 10^7$
28	$6.931 \ 10^7$	$-0.095 \ 10^7$	$8.278 \ 10^7$	$-0.096 \ 10^7$
29	$7.007 \ 10^7$	0	$8.351  10^7$	0
30	$6.928 \ 10^7$	$-0.087 \ 10^7$	$8.291  10^7$	$-0.086 \ 10^7$
31	$6.992 \ 10^7$	0	$8.338 \ 10^7$	0
32	$6.972 \ 10^7$	$-0.0001 \ 10^7$	$8.328 \ 10^7$	$-0.0002 \ 10^7$
33	$6.999 \ 10^7$	0	$8.346 \ 10^7$	0
34	$7.033 \ 10^7$	0	$8.379  10^7$	0
35	$6.928 \ 10^7$	$-0.101 \ 10^7$	$8.284 \ 10^7$	$-0.102 \ 10^7$

Table 2: One year credit Var, one year credit ES and their sensitivity

## A Proof of Proposition 1.

#### A.1 First derivative of Value at Risk.

We only provide derivation of  $\partial_{\varepsilon} VaR_3$ . The formula for  $\partial_{\varepsilon} VaR_2$  can be obtained from  $\partial_{\varepsilon} VaR_3$  by simply considering a sequence of random variables  $(Y_n)$  which tends to zero almost surely and by applying a limit theorem (such as the theorem of dominated convergence). The expression for  $\partial_{\varepsilon} VaR_1$  is a direct consequence of Property 1 in GLS (2000).

We may rewrite the equation defining  $VaR_3$  as:

$$\begin{split} \alpha &= P[L_3 > VaR_3] \\ &= P[X - \varepsilon Z > VaR_3, D = 0] \\ &+ P[X - \varepsilon Z > VaR_3, Y + \varepsilon Z \le 0, D = 1] \\ &+ P[X + Y > VaR_3, Y + \varepsilon Z > 0, D = 1] \\ &\equiv T_1 P[D = 0] + T_2 P[D = 1] + T_3 P[D = 1], \end{split}$$

where

$$T_{1} = \int \left[ \int \left\{ \int_{VaR_{3}+\epsilon z}^{+\infty} f_{(X,Y,Z)|0}(x,y,z) \, dx \right\} \, dy \right] \, dz,$$
  

$$T_{2} = \int \left[ \int_{-\infty}^{-\epsilon z} \left\{ \int_{VaR_{3}+\epsilon z}^{+\infty} f_{(X,Y,Z)|1}(x,y,z) \, dx \right\} \, dy \right] \, dz,$$
  

$$T_{3} = \int \left[ \int_{-\epsilon z}^{+\infty} \left\{ \int_{VaR_{3}-y}^{+\infty} f_{(X,Y,Z)|1}(x,y,z) \, dx \right\} \, dy \right] \, dz,$$

denoting by  $f_{U|1}$ , resp.  $f_{U|0}$ , the density of any random vector U with respects to the Lebesgue measure, conditionally to D = 1, resp. D = 0. To ease reading, set p = P[D = 1]. A simple calculation provides

$$\begin{aligned} \frac{\partial T_{1}}{\partial \varepsilon} &= -\int \left(\partial_{\varepsilon} VaR_{3} + z\right) \left\{ \int f_{(X,Y,Z)|0} (VaR_{3} + \varepsilon z, y, z) \, dy \right\} \, dz, \\ \frac{\partial T_{2}}{\partial \varepsilon} &= -\int z \left\{ \int_{VaR_{3} + \varepsilon z}^{+\infty} f_{(X,Y,Z)|1}(x, -\varepsilon z, z) \, dx \right\} \, dz \\ &- \int \left(\partial_{\varepsilon} VaR_{3} + z\right) \left\{ \int_{-\infty}^{-\varepsilon z} f_{(X,Y,Z)|1} (VaR_{3} + \varepsilon z, y, z) \, dy \right\} \, dz, \\ \frac{\partial T_{3}}{\partial \varepsilon} &= \int z \left\{ \int_{VaR_{3} + \varepsilon z}^{+\infty} f_{(X,Y,Z)|1}(x, -\varepsilon z, z) \, dx \right\} \, dz \\ &- \partial_{\varepsilon} VaR_{3} \int \left\{ \int_{-\varepsilon z}^{+\infty} f_{(X,Y,Z)|1} (VaR_{3} - y, y, z) \, dy \right\} \, dz. \end{aligned}$$

This yields

Clearly,

$$f_{(L_3,Z)|0}(l,z) = f_{(X,Z)|0}(l+\varepsilon z,z),$$
(25)

and

$$f_{(L_3,Y,Z)|1}(l,y,z) = f_{(X,Y,Z)|1}(l+\varepsilon z,y,z)\mathbf{1}\{y+\varepsilon z<0\} + f_{(X,Y,Z)|1}(l-y,y,z)\mathbf{1}\{y+\varepsilon z\ge0\}.$$
(26)

Thus, we deduce

$$\begin{array}{lcl} 0 &=& -\partial_{\varepsilon} VaR_{3} \\ & & \times \int \left\{ (1-p) \int f_{(L_{3},Y,Z)|0}(VaR_{3},y,z) \, dy \right. \\ & & + p \int_{-\infty}^{-\varepsilon z} f_{(L_{3},Y,Z)|1}(VaR_{3},y,z) \, dy + p \int_{-\varepsilon z}^{+\infty} f_{(L_{3},Y,Z)|1}(VaR_{3},y,z) \, dy \right\} \, dz \\ & & + \int z \left\{ -(1-p) \int f_{(L_{3},Y,Z)|0}(VaR_{3},y,z) \, dy - p \int_{-\infty}^{-\varepsilon z} f_{(L_{3},Y,Z)|1}(VaR_{3},y,z) \, dy \right\} \, dz \\ & = & -\partial_{\varepsilon} VaR_{3} \int \int f_{(L_{3},Y,Z)}(VaR_{3},y,z) \, dy \, dz + \int z \left\{ -(1-p) \int f_{(L_{3},Y,Z)|0}(VaR_{3},y,z) \, dy \right\} \, dz \\ & = & -\partial_{\varepsilon} VaR_{3}.f_{L_{3}}(VaR_{3}) - (1-p) \int \int z f_{(L_{3},Y,Z)|0,L_{3}=VaR_{3}}(VaR_{3},y,z) \, dy \, dz.f_{L_{3}|0}(VaR_{3}) \\ & - p \int_{-\infty}^{-\varepsilon z} z f_{(L_{3},Y,Z)|1,L_{3}=VaR_{3}}(VaR_{3},y,z) \, dy \, dz f_{L_{3}|1}(VaR_{3}). \end{array}$$

We get

$$\partial_{\varepsilon} VaR_{3} = \frac{(-p)f_{L_{3}|1}(VaR_{3})}{f_{L_{3}}(VaR_{3})} \int_{-\infty}^{-\varepsilon z} zf_{(L_{3},Y,Z)|1,L_{3}=VaR}(VaR_{3},y,z) \, dy \, dz \\ -\frac{(1-p)f_{L_{3}|0}(VaR_{3})}{f_{L_{3}}(VaR_{3})} \int_{-\infty}^{-\varepsilon z} zf_{(L_{3},Y,Z)|1,L_{3}=VaR}(VaR_{3},y,z) \, dy \, dz, \quad (27)$$

and Bayes' rule gives the stated result.  $\Box$ 

#### A.2 First derivative of Expected Shortfall.

The expression for  $\partial_{\varepsilon} ES_1$  obtains directly from Proposition 1 in Scaillet (2000). Hence we only provide derivation of  $\partial_{\varepsilon} ES_3$  since  $\partial_{\varepsilon} ES_2$  can be deduced from  $\partial_{\varepsilon} ES_3$ .

$$\begin{split} ES_3 &= E[L_3|L_3 > VaR_3] = \frac{1}{\alpha} E[L_3 \mathbf{1}\{L_3 > VaR_3)\}] \\ &= \frac{(1-p)}{\alpha} E[L_3 \mathbf{1}\{L_3 > VaR_3\}|D = 0] + \frac{p}{\alpha} E[L_3 \mathbf{1}\{L_3 > VaR_3, Y + \varepsilon Z \ge 0\}|D = 1] \\ &+ \frac{p}{\alpha} E[L_3 \mathbf{1}\{L_3 > VaR_3, Y + \varepsilon Z < 0\}|D = 1] \\ &\equiv \frac{(1-p)}{\alpha} \tilde{T}_1 + \frac{p}{\alpha} \tilde{T}_2 + \frac{p}{\alpha} \tilde{T}_3. \end{split}$$

By direct differentiation, we get

$$\begin{split} \partial_{\varepsilon} \tilde{T}_{1} &= -\int z \mathbf{1} \{ x - \varepsilon z > VaR_{3} \} f_{(X,Z)|0}(x,z) \, dx \, dz - VaR_{3} \partial_{\varepsilon} VaR_{3} \int f_{(X,Z)|0}(VaR_{3} + \varepsilon z, z) \, dz \\ &- VaR_{3} \int z f_{(X,Z)|0}(VaR_{3} + \varepsilon z, z) \, dz \\ &= -\int z \mathbf{1} \{ l > VaR_{3} \} f_{(L_{3},Z)|0}(l,z) \, dl \, dz - VaR_{3} \partial_{\varepsilon} VaR_{3} \int f_{(L_{3},Z)|0}(VaR_{3},z) \, dz \\ &- VaR_{3} \int z f_{(L_{3},Z)|0}(VaR_{3},z) \, dz. \end{split}$$

Moreover,

$$\begin{aligned} \partial_{\varepsilon} \tilde{T}_{2} &= -VaR_{3}\partial_{\varepsilon}VaR_{3}\int \mathbf{1}\{y+\varepsilon z>0\}f_{(X,Y,Z)|1}(VaR_{3}-y,y,z)\,dy\,dz \\ &+ \int (x-\varepsilon z)z\mathbf{1}\{x-\varepsilon z>VaR_{3}\}f_{(X,Y,Z)|1}(x,-\varepsilon z,z)\,dx\,dz, \end{aligned}$$

and

$$\begin{array}{ll} \partial_{\varepsilon}\tilde{T}_{3} &=& -\int z\mathbf{1}\{x-\varepsilon z > VaR_{3}, y+\varepsilon z < 0\}f_{(X,Y,Z)|1}(x,y,z)\,dx\,dy\,dz\\ &\quad -VaR_{3}\int (\partial_{\varepsilon}VaR_{3}+z)\mathbf{1}\{y+\varepsilon z < 0\}f_{(X,Y,Z)|1}(VaR_{3}+\varepsilon z,y,z)\,dy\,dz\\ &\quad -\int (x-\varepsilon z)z\mathbf{1}\{x-\varepsilon z > VaR_{3}\}f_{(X,Y,Z)|1}(x,-\varepsilon z,z)\,dx\,dz. \end{array}$$

Note that the last term of  $\partial_{\varepsilon} \tilde{T}_2$  is the opposite of the last term of  $\partial_{\varepsilon} \tilde{T}_3$ . After summing all terms, we get

$$\begin{split} \alpha \partial_{\varepsilon} ES_3 &= -(1-p) \int z \mathbf{1} \{l > VaR_3\} f_{(L_3,Z)|0}(l,z) \, dl \, dz \\ &-(1-p) VaR_3 \partial_{\varepsilon} VaR_3 \int f_{(L_3,Z)|0}(VaR_3,z) \, dz \\ &-(1-p) VaR_3 \int z f_{(L_3,Z)|0}(VaR_3,z) \, dz \\ &-p VaR_3 \partial_{\varepsilon} VaR_3 \int \mathbf{1} \{y + \varepsilon z > 0\} f_{(X,Y,Z)|1}(VaR_3 - y, y, z) \, dy \, dz \\ &-p \int z \mathbf{1} \{x - \varepsilon z > VaR_3, y + \varepsilon z < 0\} f_{(X,Y,Z)|1}(x, y, z) \, dx \, dy \, dz \\ &-p VaR_3 \partial_{\varepsilon} VaR_3 \int \mathbf{1} \{y + \varepsilon z < 0\} f_{(X,Y,Z)|1}(VaR_3 + \varepsilon z, y, z) \, dy \, dz \\ &-p VaR_3 \int z \mathbf{1} \{y + \varepsilon z < 0\} f_{(X,Y,Z)|1}(VaR_3 + \varepsilon z, y, z) \, dy \, dz \\ &= S_1 + \ldots + S_7. \end{split}$$

Note that we get by (26):

$$S_2 + S_4 + S_6 = -VaR_3\partial_{\varepsilon}VaR_3f_{L_3}(VaR_3),$$

which simplifies with

$$S_3 + S_7 = VaR_3\partial_{\varepsilon}VaR_3f_{L_3}(VaR_3),$$

because of (27). Thus

$$\begin{split} \alpha \partial_{\varepsilon} ES_3 &= -(1-p) \int \mathbf{1}\{l > VaR_3\} z f_{(L_3,Z)|0}(l,z) \, dl \, dz \\ &- p \int z \mathbf{1}\{l > VaR_3, y + \varepsilon z < 0\} f_{(L_3,Y,Z)|1}(l,y,z) \, dl \, dy \, dz \\ &= -pE[Z\mathbf{1}\{L_3 > VaR_3, Y + \varepsilon Z < 0\}|D = 1, L_3 > VaR_3]P[L_3 > VaR_3|D = 1] \\ &- (1-p)E[Z|D = 0, L_3 > VaR_3]P[L_3 > VaR_3|D = 0] \\ &= -E[Z\mathbf{1}\{Y + \varepsilon Z < 0\}|D = 1, L_3 > VaR_3]P[D = 1|L_3 > VaR_3]\alpha \\ &- E[Z|D = 0, L_3 > VaR_3]P[D = 0|L_3 > VaR_3]\alpha, \end{split}$$

proving the result.  $\square$ 

# **B** Proof of Proposition 2.

#### B.1 First derivative of Value at Risk.

Again we only need to explicit  $\partial_{\varepsilon} VaR_5$ . We may rewrite the equation defining  $VaR_5$  as:

$$\begin{aligned} \alpha &= P[L_5 > VaR_5] \\ &= (1-p)P[X > VaR_5 | D = 0] \\ &\quad + pP[X > VaR_5, Y + \varepsilon Z < 0 | D = 1] \\ &\quad + pP[X + Y + \varepsilon Z > VaR_5, Y + \varepsilon Z > 0 | D = 1] \\ &\equiv (1-p)\mathcal{T}_1 + p\mathcal{T}_2 + p\mathcal{T}_3, \end{aligned}$$

where

$$\begin{aligned} \mathcal{T}_{1} &= \int \mathbf{1}\{x > VaR_{5}\}f_{X|0}(x) \, dx, \\ \mathcal{T}_{2} &= \int \mathbf{1}\{x > VaR_{5}, y + \varepsilon z < 0\}f_{(X,Y,Z)|1}(x,y,z) \, dx \, dy \, dz, \\ \mathcal{T}_{3} &= \int \mathbf{1}\{x + y + \varepsilon z > VaR_{5}, y + \varepsilon z > 0\}f_{(X,Y,Z)|1}(x,y,z) \, dx \, dy \, dz. \end{aligned}$$

A simple computation yields

$$\begin{array}{ll} 0 &=& -(1-p)f_{X|0}(VaR_5)\partial_{\varepsilon}VaR_5 - p\partial_{\varepsilon}VaR_5 \int \mathbf{1}\{y + \varepsilon z < 0\}f_{(X,Y,Z)|1}(VaR_5,y,z)\,dy\,dz\\ &\quad -p\int z\mathbf{1}\{x > VaR_5\}f_{(X,Y,Z)|1}(x,-\varepsilon z,z)\,dx\,dz\\ &\quad -p\int (\partial_{\varepsilon}VaR_5 - z)\mathbf{1}\{y + \varepsilon z > 0\}f_{(X,Y,Z)|1}(VaR_5 - y - \varepsilon z,y,z)\,dy\,dz\\ &\quad +p\int z\mathbf{1}\{x > VaR_5\}f_{(X,Y,Z)|1}(x,-\varepsilon z,z)\,dx\,dz. \end{array}$$

Clearly, we have

$$f_{(L_5,Y,Z)|0}(l,y,z) = f_{(X,Y,Z)|0}(l,y,z),$$
(28)

and

$$f_{(L_5,Y,Z)|1}(l,y,z) = \mathbf{1}\{y + \varepsilon z > 0\} f_{(X,Y,Z)|1}(l-y-\varepsilon z,y,z) + \mathbf{1}\{y + \varepsilon z < 0\} f_{(X,Y,Z)|1}(l,y,z).$$
(29)

Thus, we can rewrite

$$0 = -\partial_{\varepsilon} VaR_{5} f_{L_{5}}(VaR_{5}) + p \int z \mathbf{1} \{y + \varepsilon z > 0\} f_{(L_{5}, Y, Z)|1}(VaR_{5}, y, z) \, dy \, dz.$$
(30)

We deduce

$$\partial_{\varepsilon} VaR_5 \frac{f_{L_5}(VaR_5)}{f_{L_5|1}(VaR_5)} = P[D=1]E[Z\mathbf{1}\{Y + \varepsilon Z > 0\}|D=1, L_5 = VaR_5],$$

and finally

$$\partial_{\varepsilon} VaR_5 = P[D = 1|L_5 = VaR_5]E[Z\mathbf{1}\{Y + \varepsilon Z > 0\}|D = 1, L_5 = VaR_5],$$

by Bayes' rule.  $\Box$ 

# B.2 First derivative of Expected Shortfall.

We only provide derivation of  $\partial_{\varepsilon} ES_5$ . As previously,

$$\begin{split} ES_5 &= E[L_5|L_5 > VaR_5] = \frac{1}{\alpha} E[L_5 \mathbf{1}\{L_5 > VaR_5\}] \\ &= \frac{1-p}{\alpha} E[L_5 \mathbf{1}\{L_5 > VaR_5\}|D = 0] + \frac{p}{\alpha} E[L_5 \mathbf{1}\{L_5 > VaR_5, Y + \varepsilon Z < 0\}|D = 1] \\ &\quad + \frac{p}{\alpha} E[L_5 \mathbf{1}\{L_5 > VaR_5, Y + \varepsilon Z \ge 0\}|D = 1] \\ &= \frac{1-p}{\alpha} \int x \mathbf{1}\{x \ge VaR_5\} f_{X|0}(x) \, dx \\ &\quad + \frac{p}{\alpha} \int x \mathbf{1}\{x \ge VaR_5, y + \varepsilon z < 0\} f_{(X,Y,Z)|1}(x, y, z) \, dx \, dy \, dz \\ &\quad + \frac{p}{\alpha} \int (x + y + \varepsilon z) \mathbf{1}\{(x + y + \varepsilon z) \ge VaR_5, y + \varepsilon z \ge 0\} f_{(X,Y,Z)|1}(x, y, z) \, dx \, dy \, dz \\ &\equiv \frac{1-p}{\alpha} \tilde{\mathcal{T}}_1 + \frac{p}{\alpha} \tilde{\mathcal{T}}_2 + \frac{p}{\alpha} \tilde{\mathcal{T}}_3. \end{split}$$

Here,

$$\begin{array}{lll} \partial_{\varepsilon}\tilde{\mathcal{T}_{1}} &=& -VaR_{5}\partial_{\varepsilon}VaR_{5}f_{X|0}(VaR_{5}),\\ \partial_{\varepsilon}\tilde{\mathcal{T}_{2}} &=& -VaR_{5}\partial_{\varepsilon}VaR_{5}\int\mathbf{1}\{y+\varepsilon z<0\}f_{(X,Y,Z)|1}(VaR_{5},y,z)\,dy\,dz\\ && -\int zx\mathbf{1}\{x>VaR_{5}\}f_{(X,Y,Z)|1}(x,-\varepsilon z,z)\,dx\,dz,\\ \partial_{\varepsilon}\tilde{\mathcal{T}_{3}} &=& \int z\mathbf{1}\{x+y+\varepsilon z>VaR_{5},y+\varepsilon z\ge0\}f_{(X,Y,Z)|1}(x,y,z)\,dx\,dy\,dz\\ && -\int (\partial_{\varepsilon}VaR_{5}-z)VaR_{5}\mathbf{1}\{y+\varepsilon z>0\}f_{(X,Y,Z)|1}(VaR_{5}-y-\varepsilon z,y,z)\,dy\,dz\\ && +\int zx\mathbf{1}\{x>VaR_{5}\}f_{(X,Y,Z)|1}(x,-\varepsilon z,z)\,dx\,dz. \end{array}$$

We get by recalling Equation (30):

$$\begin{split} \alpha \partial_{\varepsilon} ES_5 &= -VaR_5 \partial_{\varepsilon} VaR_5 f_{L_5}(VaR_5) \\ &+ p \int z \mathbf{1} \{l > VaR_5, y + \varepsilon z > 0\} f_{(L_5,Y,Z)|1}(l,y,z) \, dl \, dy \, dz \\ &+ p VaR_5 \int z \mathbf{1} \{y + \varepsilon z > 0\} f_{(L_5,Y,Z)|1}(VaR_5,y,z) \, dl \, dy \, dz \\ &= p \int z \mathbf{1} \{l > VaR_5, y + \varepsilon z > 0\} f_{(L_5,Y,Z)|1}(l,y,z) \, dl \, dy \, dz. \end{split}$$

Thus, by applying Bayes' rule, we get

$$\partial_{\varepsilon} ES_5 = \frac{P[D=1]}{P[L_5 > VaR_5]} \int z \mathbf{1} \{l > VaR_5, y + \varepsilon z > 0\} f_{(L_5, Y, Z)|1}(l, y, z) \, dl \, dy \, dz$$
  
=  $P[D=1|L_5 > VaR_5] E[Z \mathbf{1} \{Y + \varepsilon Z > 0\} | L_5 > VaR_5, D = 1],$ 

which proves the result.  $\Box$ 

## C Proof of Proposition 3.

Since the calculations are very similar to previous ones, we prove the results for  $L'_5$  only and we omit unnecessary details.

#### C.1 First derivative of Value at Risk.

We may rewrite the equation defining  $VaR'_5$  as:

$$\begin{split} \alpha &= P[L_5' > VaR_5'] &= (1-p)P[X > VaR_5'|D=0] + pP[X > VaR_5', Y + \varepsilon Z < 0|D=1] \\ &+ pP[X + Y + \varepsilon Z > VaR_5', Y + \varepsilon Z \in [0, M]|D=1] \\ &+ pP[X + M > VaR_5', Y + \varepsilon Z > M|D=1]. \end{split}$$

Simple computations and some simplifications yield

$$\begin{array}{ll} 0 &=& -(1-p)f_{X|0}(VaR_5')\partial_{\varepsilon}VaR_5' - p\partial_{\varepsilon}VaR_5' \int \mathbf{1}\{y + \varepsilon z < 0\}f_{(X,Y,Z)|1}(VaR_5',y,z)\,dy\,dz\\ && -p\int(\partial_{\varepsilon}VaR_5'-z)\mathbf{1}\{y + \varepsilon z \in [0,M]\}f_{(X,Y,Z)|1}(VaR_5'-y - \varepsilon z,y,z)\,dy\,dz\\ && -p\partial_{\varepsilon}VaR_5'\int \mathbf{1}\{y + \varepsilon z > M\}f_{(X,Y,Z)|1}(VaR_5'-M,y,z)\,dy\,dz. \end{array}$$

Clearly, we have

$$\begin{split} f_{(L'_{5},Y,Z)|1}(l,y,z) &= \mathbf{1}\{y + \varepsilon z \in [0,M]\} f_{(X,Y,Z)|1}(l-y-\varepsilon z,y,z) \\ &+ \mathbf{1}\{y + \varepsilon z < 0\} f_{(X,Y,Z)|1}(l,y,z) \\ &+ \mathbf{1}\{y + \varepsilon z > M\} f_{(X,Y,Z)|1}(l-M,y,z). \end{split}$$

Thus, we can rewrite

$$0 = -\partial_{\varepsilon} VaR'_{5} f_{L'_{5}}(VaR'_{5}) + p \int z \mathbf{1}\{y + \varepsilon z \in [0, M]\} f_{(L'_{5}, Y, Z)|1}(VaR'_{5}, y, z) \, dy \, dz, \quad (31)$$

so the stated result.  $\square$ 

# C.2 First derivative of Expected Shortfall.

As previously,

$$\begin{split} ES'_{5} &= \frac{1-p}{\alpha} E[L'_{5}\mathbf{1}\{L'_{5} > VaR'_{5}\}|D = 0] + \frac{p}{\alpha} E[L'_{5}\mathbf{1}\{L'_{5} > VaR'_{5}, Y + \varepsilon Z < 0\}|D = 1] \\ &\quad + \frac{p}{\alpha} E[L'_{5}\mathbf{1}\{L'_{5} > VaR'_{5}, Y + \varepsilon Z \in [0, M]\}|D = 1] \\ &\quad + \frac{p}{\alpha} E[L'_{5}\mathbf{1}\{L'_{5} > VaR'_{5}, Y + \varepsilon Z > M\}|D = 1] \\ &= \frac{1-p}{\alpha} \int x\mathbf{1}\{x \ge VaR'_{5}\}f_{X|0}(x) \, dx \\ &\quad + \frac{p}{\alpha} \int x\mathbf{1}\{x \ge VaR'_{5}, y + \varepsilon z < 0\}f_{(X,Y,Z)|1}(x, y, z) \, dx \, dy \, dz \\ &\quad + \frac{p}{\alpha} \int (x + y + \varepsilon z)\mathbf{1}\{(x + y + \varepsilon z) \ge VaR'_{5}, y + \varepsilon z \in [0, M]\}f_{(X,Y,Z)|1}(x, y, z) \, dx \, dy \, dz \\ &\quad + \frac{p}{\alpha} \int (x + M)\mathbf{1}\{(x + M) \ge VaR'_{5}, y + \varepsilon z > M\}f_{(X,Y,Z)|1}(x, y, z) \, dx \, dy \, dz \\ &\equiv \frac{1-p}{\alpha}\tilde{T}'_{1} + \frac{p}{\alpha}\tilde{T}'_{2} + \frac{p}{\alpha}\tilde{T}'_{3} + \frac{p}{\alpha}\tilde{T}'_{4}. \end{split}$$

Here,

$$\begin{array}{lll} \partial_{\varepsilon}\tilde{T}'_{1} &=& -VaR'_{5}\partial_{\varepsilon}VaR'_{5}f_{X|0}(VaR'_{5}),\\ \partial_{\varepsilon}\tilde{T}'_{2} &=& -VaR'_{5}\partial_{\varepsilon}VaR'_{5}\int\mathbf{1}\{y+\varepsilon z<0\}f_{(X,Y,Z)|1}(VaR'_{5},y,z)\,dy\,dz\\ && -\int zx\mathbf{1}\{x>VaR'_{5}\}f_{(X,Y,Z)|1}(x,-\varepsilon z,z)\,dx\,dz,\\ \partial_{\varepsilon}\tilde{T}'_{3} &=& \int z\mathbf{1}\{x+y+\varepsilon z>VaR'_{5},y+\varepsilon z\in[0,M]\}f_{(X,Y,Z)|1}(x,y,z)\,dx\,dy\,dz\\ && -\int (\partial_{\varepsilon}VaR'_{5}-z)VaR'_{5}\mathbf{1}\{y+\varepsilon z\in[0,M]\}f_{(X,Y,Z)|1}(VaR'_{5}-y-\varepsilon z,y,z)\,dy\,dz\\ && +\int zx\mathbf{1}\{x>VaR'_{5}\}f_{(X,Y,Z)|1}(x,-\varepsilon z,z)\,dx\,dz\\ && -\int z(x+M)\mathbf{1}\{x+M>VaR'_{5}\}f_{(X,Y,Z)|1}(x,M-\varepsilon z,z)\,dx\,dz,\\ \partial_{\varepsilon}\tilde{T}'_{4} &=& -VaR'_{5}\partial_{\varepsilon}VaR'_{5}\int\mathbf{1}\{y+\varepsilon z>M\}f_{(X,Y,Z)|1}(VaR'_{5}-M,y,z)\,dy\,dz\\ && +\int z(x+M)\mathbf{1}\{x+M>VaR'_{5}\}f_{(X,Y,Z)|1}(x,M-\varepsilon z,z)\,dx\,dz. \end{array}$$

After some simplifications and invoking Equation (31), we get

$$\begin{split} \alpha \partial_{\varepsilon} ES'_{5} &= -VaR'_{5} \partial_{\varepsilon} VaR'_{5} f_{L'_{5}}(VaR'_{5}) \\ &+ p \int z \mathbf{1}\{l > VaR'_{5}, y + \varepsilon z \in [0, M]\} f_{(L'_{5}, Y, Z)|1}(l, y, z) \, dl \, dy \, dz \\ &+ p VaR'_{5} \int z \mathbf{1}\{y + \varepsilon z \in [0, M]\} f_{(L'_{5}, Y, Z)|1}(VaR'_{5}, y, z) \, dl \, dy \, dz \\ &= p \int z \mathbf{1}\{l > VaR'_{5}, y + \varepsilon z \in [0, M]\} f_{(L'_{5}, Y, Z)|1}(l, y, z) \, dl \, dy \, dz, \end{split}$$

and finally the result.  $\Box$ 

# **D** Sensitivity of *VaR* and *ES* under the Gaussian assumption.

Let us first recall a useful lemma:

**Lemma 5.** The law of  $E_1$  conditionally to  $E_2 = e$  is Gaussian with mean

$$m_{1|2}(e) = m_1 + \frac{\rho \sigma_1}{\sigma_2}(e - m_2),$$

 $and \ variance$ 

$$\sigma_{1|2}^2 = \sigma_1^2 (1 - \rho^2).$$

Invoking (13), the sensitivity of  $L_U$ 's Value-At-Risk is

$$\partial_{\varepsilon} VaR_U = E[E_2 \mathbf{1}\{E_2 > 0, D = 1\} | L_U = v],$$

where v is the VaR level. First, let us calculate the density of  $L_U$ . Actually,  $L_U$  equals 0 with the probability

$$P(L_U = 0) = 1 - p + pP(E_1 < 0, E_2 < 0).$$

Otherwise,  $L_U$  has a density with respects to the Lebesgue measure on  $\mathbb{R}^+$ .

More specifically, consider a measurable real function  $\psi$ . Then

$$\begin{split} E[\psi(L_U)] &= \psi(0)\{1 - p + pP(E_1 < 0, E_2 < 0)\} \\ &+ p \int \psi(e) \mathbf{1}\{e > 0\} P(E_2 < 0|E_1 = e) f_{E_1}(e) \, de \\ &+ p \int \psi(\varepsilon e) \mathbf{1}\{e > 0\} P(E_1 < 0|E_2 = e) f_{E_2}(e) \, de \\ &+ p \int \psi(l) \mathbf{1}\{l > 0, e > 0\} f_{(\overline{L}, E_2)}(l, e) \, de \, dl, \end{split}$$

where we have set  $\overline{L} = E_1 + \varepsilon E_2$ . By applying successively Lemma 5, we get

$$P(E_2 < 0|E_1 = e) = \Phi\left(-\left(m_2 + \frac{\rho\sigma_2}{\sigma_1}(e - m_1)\right) \cdot \frac{1}{\sigma_2\sqrt{1 - \rho^2}}\right),$$
$$P(E_1 < 0|E_2 = e) = \Phi\left(-\left(m_1 + \frac{\rho\sigma_1}{\sigma_2}(e - m_2)\right) \cdot \frac{1}{\sigma_1\sqrt{1 - \rho^2}}\right).$$

Moreover, the random vector  $(\overline{L}, E_2)$  is Gaussian with mean  $m = (m_1 + \varepsilon m_2, m_2)'$  and covariance matrix

$$\overline{\Omega} = \begin{bmatrix} \tau^2 & \nu \tau \sigma_2 \\ \\ \nu \tau \sigma_2 & \sigma_2^2, \end{bmatrix}.$$

with

$$\tau^{2} = \sigma_{1}^{2} + \varepsilon^{2} \sigma_{2}^{2} + 2\varepsilon \rho \sigma_{1} \sigma_{2},$$
$$\nu \tau \sigma_{2} = \rho \sigma_{1} \sigma_{2} + \varepsilon \sigma_{2}^{2}.$$

Thus, by applying Lemma 5 and the symmetry of the Gaussian distribution, we get

$$P(E_2 > 0 | \overline{L} = l) = \Phi\left(\frac{1}{\sigma_2 \sqrt{1 - \nu^2}} [m_2 + \frac{\nu \sigma_2}{\tau} (l - m_1 - \varepsilon m_2)]\right).$$

The density of  $L_U$  at l > 0 is then

$$f_{L_U}(l) = p \frac{P(E_2 < 0|E_1 = l)}{\sigma_1} \phi\left(\frac{(l - m_1)}{\sigma_1}\right) + p \frac{P(E_1 < 0|E_2 = l/\varepsilon)}{\varepsilon \sigma_2} \phi\left(\frac{(l - m_2\varepsilon)}{\varepsilon \sigma_2}\right) + p \frac{P(E_2 > 0|\overline{L} = l)}{\tau} \phi\left(\frac{(l - m_1 - \varepsilon m_2)}{\tau}\right).$$
(32)

Moreover, for every v > 0 such that  $f_{L_U}(v) > 0$ ,

$$\partial_{\varepsilon} VaR_{U} = E[E_{2}\mathbf{1}\{E_{2} > 0, D = 1\}|L_{U} = v]$$

$$= E[E_{2}\mathbf{1}\{E_{1} > 0, E_{2} > 0, D = 1\}|L_{U} = v] + E[E_{2}\mathbf{1}\{E_{1} < 0, E_{2} > 0, D = 1\}|L_{U} = v]$$

$$= E[E_{2}\mathbf{1}\{E_{1} > 0, E_{2} > 0, D = 1\}|L_{U} = v] + E[\frac{L_{U}}{\varepsilon}\mathbf{1}\{E_{1} < 0, L_{U} > 0, D = 1\}|L_{U} = v]$$

$$= \frac{p}{f_{L_{U}}(v)} \left[\int e\mathbf{1}\{v - \varepsilon e > 0, e > 0\}f_{(E_{1}, E_{2})}(v - \varepsilon e, e) de$$

$$+ \frac{v}{\varepsilon}\int \mathbf{1}\{e < 0\}f_{(E_{1}, E_{2})}(e, \frac{v}{\varepsilon}) de\right].$$
(33)

For convenience, denote  $\tilde{e} = e - m_2$  and  $\tilde{v} = v - m_1 - \varepsilon m_2$ . Let us compute

$$\int e\mathbf{1}\{v - \varepsilon e > 0, e > 0\} f_{(E_1, E_2)}(v - \varepsilon e, e) de$$

$$= \int e\mathbf{1}\{v - \varepsilon e > 0, e > 0\} \exp\left(-\frac{\sigma_2^2 [\tilde{v} - \varepsilon \tilde{e}]^2 + \sigma_1^2 \tilde{e}^2 - 2\rho\sigma_1\sigma_2[\tilde{v} - \varepsilon \tilde{e}]\tilde{e}}{2\sigma_1^2 \sigma_2^2 (1 - \rho^2)}\right) \frac{de}{2\pi\sigma_1\sigma_2\sqrt{1 - \rho^2}}$$

$$= \int e\mathbf{1}\{v - \varepsilon e > 0, e > 0\} \exp\left(-\frac{[\tau \tilde{e} - \nu \tilde{v}\sigma_2]^2 + (1 - \nu^2)\sigma_2^2 \tilde{v}^2}{2\sigma_1^2 \sigma_2^2 (1 - \rho^2)}\right) \frac{de}{2\pi\sigma_1\sigma_2\sqrt{1 - \rho^2}}$$

$$= \phi\left(\frac{\tilde{v}\sqrt{1 - \nu^2}}{\sigma_1\sqrt{1 - \rho^2}}\right) \int e\mathbf{1}\{v - \varepsilon e > 0, e > 0\} \phi\left(\frac{\tau \tilde{e} - \nu\sigma_2 \tilde{v}}{\sigma_1\sigma_2\sqrt{1 - \rho^2}}\right) \frac{de}{\sigma_1\sigma_2\sqrt{1 - \rho^2}}$$

$$= \frac{1}{\tau^2} \phi\left(\frac{\tilde{v}\sqrt{1 - \nu^2}}{\sigma_1\sqrt{1 - \rho^2}}\right) \cdot \left[\sigma_1\sigma_2\sqrt{1 - \rho^2} \int_c^d t\phi(t) dt + (m_2\tau + \nu\sigma_2 \tilde{v}) \int_c^d \phi(t) dt\right]$$

$$= \frac{1}{\tau^2} \phi\left(\frac{\tilde{v}\sqrt{1 - \nu^2}}{\sigma_1\sqrt{1 - \rho^2}}\right) \cdot \left[\sigma_1\sigma_2\sqrt{1 - \rho^2} (\phi(c) - \phi(d)) + (m_2\tau + \nu\sigma_2 \tilde{v}) (\Phi(d) - \Phi(c))\right], \quad (34)$$

by setting

$$c = -\frac{m_2\tau}{\sigma_1\sigma_2\sqrt{1-\rho^2}} - \frac{\nu\tilde{v}}{\sigma_1\sqrt{1-\rho^2}}, \quad d = \left(\frac{v}{\varepsilon} - \frac{\nu\sigma_2\tilde{v}}{\tau} - m_2\right) \cdot \frac{\tau}{\sigma_1\sigma_2\sqrt{1-\rho^2}}.$$

and by denoting  $\phi$ , resp.  $\Phi$ , the pdf, resp. cdf of a standard Gaussian random variable. Moreover,

$$\int \mathbf{1}\{e < 0\} f_{(E_1, E_2)}(e, \frac{v}{\varepsilon}) de$$

$$= \phi \left(\frac{v/\varepsilon - m_2}{\sigma_2}\right) \int \mathbf{1}\{e < 0\} \exp\left(-\frac{[\sigma_2(e - m_1) - \rho\sigma_1(v/\varepsilon - m_2)]^2}{2\sigma_1^2 \sigma_2^2 (1 - \rho^2)}\right) \frac{de}{\sqrt{2\pi}\sigma_1 \sigma_2 \sqrt{1 - \rho^2}}$$

$$= \frac{1}{\sigma_2} \phi \left(\frac{v - \varepsilon m_2}{\varepsilon \sigma_2}\right) \Phi \left(-\frac{m_1}{\sigma_1 \sqrt{1 - \rho^2}} - \frac{\rho(v - \varepsilon m_2)}{\varepsilon \sigma_2 \sqrt{1 - \rho^2}}\right). \tag{35}$$

By taking together equations (33), (32), (34) and (35), we find  $\partial_{\varepsilon} VaR_U$ .

Similar calculations provide  $\partial_{\varepsilon} VaR_N$  in the netted case. Here, the loss function equals zero with the probability  $1-p+pP(E_1+\varepsilon E_2 < 0)$ . This function has a density with respect to the Lebesgue measure at every positive point v, and, with the previous notations,

$$f_{L_N}(v) = \frac{p}{\tau} \cdot \phi\left(\frac{v - m_1 - \varepsilon m_2}{\tau}\right) \mathbf{1}\{v > 0\}.$$
(36)

Moreover,

$$\partial_{\varepsilon} VaR_{N} = E[E_{2}\mathbf{1}\{D=1\}|L_{N}=v]$$

$$= \frac{p}{f_{L_{N}}(v)} \int ef_{(\overline{L},E_{2})}(v,e) de$$

$$= \frac{p}{f_{L_{N}}(v)} \int e\exp\left(-\frac{[\tau \tilde{e} - \nu \sigma_{2}\tilde{v}]^{2}}{2\sigma_{2}^{2}\tau^{2}(1-\nu^{2})}\right) \frac{de}{\sqrt{2\pi}\sigma_{2}\tau\sqrt{1-\nu^{2}}} \cdot \phi\left(\frac{\tilde{v}}{\tau}\right)$$

$$= \frac{p}{f_{L_{N}}(v)} \int \left(\sigma_{2}\tau\sqrt{1-\nu^{2}}u + \nu\sigma_{2}\tilde{v} + m_{2}\tau\right)\phi(u) \frac{du}{\tau^{2}} \cdot \phi\left(\frac{\tilde{v}}{\tau}\right)$$

$$= \frac{p}{\tau^{2}f_{L_{N}}(v)} \cdot \phi\left(\frac{\tilde{v}}{\tau}\right) [\nu\sigma_{2}\tilde{v} + \tau m_{2}]. \tag{37}$$

Equations (36) and (37) provide the sensitivity of the VaR for  $L_N$ .  $\Box$ 

Let us remark that the sensitivities of the expected shortfalls of  $L_N$  and  $L_U$  can be calculated in the same way. Unfortunately the formulas are more complicated. An extra integration with respect to v is in fact needed, and the sensitivities do not admit simple closed forms.

#### **E** Asymptotic normality of *VaR* sensitivity estimators

Let us consider an i.i.d. sample  $(e_i, l_i)_{i=1,...,n}$  of a random vector (e, l). The density of l is denoted by f. Let  $\hat{v}$  be a statistic which tends to a constant v and  $n^{1/2}(\hat{v} - v) = O_P(1)$ . Set

$$\hat{r}(v) = \frac{\sum_{i=1}^{n} e_i K_h(l_i - \hat{v})}{\sum_{i=1}^{n} K_h(l_i - \hat{v})},$$

and denote the expectation of  $e_i$  conditionally to  $l_i = v$  by r(v), and its conditional variance by

$$\mu(v) = E[e_i^2 | l_i = v] - r(v)^2.$$

#### Lemma 6. Assume

- K is an even kernel function, ∫ |t|<sup>3</sup>|K|(t) dt < ∞, lim<sub>|t|→∞</sub> |t|<sup>3</sup>K(t) = 0. It is three times continuously differentiable, K' and K'' are integrable and K''' is bounded.
- $E[|e|^p] < \infty$  for every integer p,
- $t \mapsto E[e^4|l=t]$  is bounded in a neighborhood of v,
- $nh^5 \longrightarrow 0$  and  $nh^{7/2} \longrightarrow \infty$ ,
- r and f are two times continuously differentiable,
- f and  $\mu$  are continuous and strictly positive in a neighborhood of v.

Then

$$\sqrt{nh}\left\{\hat{r}(v) - r(v)\right\} \xrightarrow[n \to \infty]{law} \mathcal{N}\left(0, \int K^2 \cdot \mu(v) / f(v)\right).$$

Obviously the limiting behavior of  $\partial_{\varepsilon} \widehat{VaR}_k$ , k = 4, 5 is a direct consequence of the previous lemma. For instance, set  $e_i = z_i \mathbf{1}\{z_i > 0, D_i = 1\}$  to get the result for  $\partial_{\varepsilon} \widehat{VaR}_4$ .

#### Proof of Lemma 6

The estimator  $\hat{r}(v)$  can be decomposed into

$$\hat{r}(v) = \frac{\sum_{i=1}^{n} e_i K_h(l_i - v)}{\sum_{i=1}^{n} K_h(l_i - v)} + R_1(v) + \hat{r}(v) R_2(v),$$
(38)

with

$$R_1(v) = \frac{n^{-1} \sum_{i=1}^n e_i \left\{ K_h(l_i - \hat{v}) - K_h(l_i - v) \right\}}{n^{-1} \sum_{i=1}^n K_h(l_i - v)},$$

and

$$R_2(v) = \frac{n^{-1} \sum_{i=1}^n \left\{ K_h(l_i - v) - K_h(l_i - \hat{v}) \right\}}{n^{-1} \sum_{i=1}^n K_h(l_i - v)}$$

By applying Corollary IV.2 in Bosq and Lecoutre (1987), we get

$$\sqrt{nh} \left\{ \frac{\sum_{i=1}^{n} e_i K_h(l_i - v)}{\sum_{i=1}^{n} K_h(l_i - v)} - E[e|l = v] \right\} \xrightarrow[n \to \infty]{law} \mathcal{N}\left(0, \int K^2 \cdot \mu(v) / f(v)\right).$$
(39)

Moreover,

$$P\left(\sqrt{nh}|R_{1}(v)| > \varepsilon\right) \leq P\left(\frac{\sqrt{nh}}{n}|\sum_{i=1}^{n} e_{i}\left\{K_{h}(l_{i}-\hat{v}) - K_{h}(l_{i}-v)\right\}| > \varepsilon f(v)/2\right) + P\left(|n^{-1}\sum_{i=1}^{n} K_{h}(l_{i}-v) - f(v)| > f(v)/2\right) \equiv P_{1} + P_{2}.$$

The second term on the right hand side tends to zero (convergence in probability of the usual kernel estimator of the density function). By a three order expansion of  $K_h$ , we get

$$P_{1} \leq P\left(\frac{\sqrt{nh}}{nh}|\sum_{i=1}^{n}e_{i}\left\{(K')_{h}(l_{i}-v).(\hat{v}-v)\right\}| > \varepsilon f(v)/6\right)$$
  
+ $P\left(\frac{\sqrt{nh}}{nh^{2}}|\sum_{i=1}^{n}e_{i}\left\{(K'')_{h}(l_{i}-v).(\hat{v}-v)^{2}\right\}| > \varepsilon f(v)/3\right)$   
+ $P\left(\frac{\sqrt{nh}}{nh^{3}}|\sum_{i=1}^{n}e_{i}\left\{(K''')_{h}(l_{i}-v^{*}).(\hat{v}-v)^{3}\right\}| > \varepsilon f(v)\right) \equiv P_{11} + P_{12} + P_{13},$ 

where  $|v^* - v| < |\hat{v} - v|$  a.e. Clearly,  $P_{13}$  is zero for *n* sufficiently large when  $nh^{7/2} \to \infty$ . Moreover, for every constant  $\eta > 0$ , there exists a constant C > 0 such that

$$P_{11} \le P(n^{1/2}|\hat{v} - v| > C) + P\left(\frac{\sqrt{h}}{nh}|\sum_{i=1}^{n} e_i\left\{(K')_h(l_i - v)\right\}| > \varepsilon f(v)/(6C)\right)$$
  
$$\le \eta + \frac{E[|e_i(K')_h(l_i - v)|]}{\sqrt{Cst.h}}$$
  
$$\le \eta + O(h^{1/2}),$$

by an integration by parts. Thus,  $P_{11}$  tends to zero when  $n \to \infty$ . Similar calculations provide the same result for  $P_{12}$ . Thus, we deduce

$$R_1(v) = o_P(1). (40)$$

We can handle  $R_2(v)$  similarly. Thus, we get the stated result by combining (38), (39) and (40).  $\Box$ 

### References

- ACERBI C. and D. TASCHE (2002). On the coherence of expected shortfall. Journal of Banking and Finance, 26, 1487-1503.
- [2] BANK FOR INTERNATIONAL SETTLEMENT (2003). OTC derivatives market activity in the second half of 2002.
- BASAK S. and A. SHAPIRO (2001). Value-at-Risk Based Risk Management: Optimal Policies and Asset Prices. *Review of Financial Studies*, December, 371-405.
- [4] BASEL COMMITTEE ON BANKING SUPERVISION (2001) The new Basel Capital Accord. Consultative document. Bank for International Settlements.
- [5] BOSQ D. and J-P. LECOUTRE (1987). *Théorie de l'estimation fonctionnelle*. Economica. Paris.
- [6] CROUHY, M., D. GALAI and R. MARK (1998). The new 1998 regulatory framework for capital adequacy :"Standardized approach" versus "internal models". In *Risk management and analysis*, edited by C. Alexander. Wiley.
- [7] CUOCO D., H. HE and S. ISSAENKO (2001). Optimal Dynamic Trading Strategies with Risk Limits, DP Wharton.
- [8] DENAULT M. (2001). Coherent Allocation of Risk Capital. Journal of Risk, 4, 7-21.
- [9] DUFFIE D. and K. SINGLETON (2003). Credit Risk. Pricing, Measurement and Management, Princeton University Press, Princeton.
- [10] FREY R. and A. MCNEIL (2002). VaR and Expected Shortfall in Portfolios of Dependent Credit Risks: Conceptual and Practical Insights. *Journal of Banking and Finance*, 26, 1317-1334.
- [11] GARMAN M. (1996). Improving on VaR. RISK, 9, 61-63.
- [12] GARMAN M. (1997). Taking VaR to Pieces. RISK, 10, 70-71.



Figure 1: Modélisation des taux de défauts mensuels sur données US

- [13] GOURIÉROUX C., J.P. LAURENT and O. SCAILLET (2000). Sensitivity Analysis of Values at Risk. *Journal of Empirical Finance*, 7, 225-245.
- [14] HALLERBACH W. (2003). Decomposing Portfolio Value-at-Risk: a General Analysis. Forthcoming in *Journal of Risk*.
- [15] JORION P. (1997). Value at Risk: The New Benchmark for Controlling Market Risk, Irwin, Chicago.
- [16] KURTH A. and D. TASCHE (2002). Credit Risk Contributions to Value-at-Risk and Expected Shortfall, DP TU München.
- [17] LEIPPOLD M., F. TROJANI and P. VANINI (2002). Equilibrium Impact of Value-at-Risk, DP University of Zurich.
- [18] MARKOWITZ H. (1952). Portfolio Selection. Journal of Finance, 7, 77-91.
- [19] MARTIN R., K. THOMPSON and C. BROWNE (2001). VaR: Who Contributes and How Much? *Risk*, August, 99-102.
- [20] MARTIN R. and T. WILDE (2002). Unsystematic credit risk? *Risk*, November, 123-128.
- [21] ROCKAFELLAR T. and S. URYASEV (2000). Optimization of Conditional Value at Risk. Journal of Risk, 2, 21-41.
- [22] ROCKAFELLAR T. and S. URYASEV (2002). Conditional Value-at-Risk for General Loss Distributions. *Journal of Banking and Finance*, 26, 1443-1471.
- [23] SCAILLET O. (2000). Nonparametric Estimation and Sensitivity Analysis of Expected Shortfall. Forthcoming in *Mathematical Finance*.
- [24] SZEGÖ G. (2002). Measures of Risk. Journal of Banking and Finance, 26, 1253-1272.
- [25] TASCHE D. (1999). Risk Contributions and Performance Measurement, DP TU München.

[26] VAN DER VAART A. (1994). Weak convergence of smoothed empirical processes. Scandinavian Journal of Statistics, 21, 501-504.